

Bath Observables with HOPS

Energy Flow in Strongly Coupled Open Quantum Systems

Valentin Boettcher, Richard Hartmann, Konstantin Beyer, Walter Strunz

Institute for Theoretical Physics, Dresden, Germany

17.08.2022



Introduction

- Motivation

- Technical Basics

Bath and Interaction Energy

- A Little (more) Theory

- Analytic Verification

Applications

- One Bath

- Energy Shovel

- Otto Cycle

- Anti-Zeno Engine

Outlook

Introduction

- Motivation

- Technical Basics

Bath and Interaction Energy

- A Little (more) Theory

- Analytic Verification

Applications

- One Bath

- Energy Shovel

- Otto Cycle

- Anti-Zeno Engine

Outlook

Introduction

Motivation

Technical Basics

Bath and Interaction Energy

A Little (more) Theory

Analytic Verification

Applications

One Bath

Energy Shovel

Otto Cycle

Anti-Zeno Engine

Outlook

Situation

Consider an open quantum system

$$H = \underbrace{H_S}_{\text{"small"}} + \underbrace{H_I}_{?} + \underbrace{H_B}_{\text{"big", simple}} \quad (1)$$

with $[H_S, H_B] = 0$.

¹Rivas, "Strong Coupling Thermodynamics of Open Quantum Systems"; Talkner and Hnggi, "Colloquium: Statistical mechanics and thermodynamics at strong coupling: Quantum and classical".

Situation

Consider an open quantum system

$$H = \underbrace{H_S}_{\text{"small"}} + \underbrace{H_I}_{?} + \underbrace{H_B}_{\text{"big", simple}} \quad (1)$$

with $[H_S, H_B] = 0$.

- ▶ weak coupling $H_I \approx 0$ thermodynamics of open systems are somewhat understood¹

¹Rivas, "Strong Coupling Thermodynamics of Open Quantum Systems"; Talkner and Hnggi, "Colloquium: Statistical mechanics and thermodynamics at strong coupling: Quantum and classical".

Situation

Consider an open quantum system

$$H = \underbrace{H_S}_{\text{"small"}} + \underbrace{H_I}_{?} + \underbrace{H_B}_{\text{"big", simple}} \quad (1)$$

with $[H_S, H_B] = 0$.

- ▶ weak coupling $H_I \approx 0$ thermodynamics of open systems are somewhat understood¹
- ▶ strong coupling: $\langle H_I \rangle \sim \langle H_S \rangle \implies$ we can't neglect the interaction \implies thermodynamics?

¹Rivas, "Strong Coupling Thermodynamics of Open Quantum Systems"; Talkner and Hnggi, "Colloquium: Statistical mechanics and thermodynamics at strong coupling: Quantum and classical".

Situation

Consider an open quantum system

$$H = \underbrace{H_S}_{\text{"small"}} + \underbrace{H_I}_{?} + \underbrace{H_B}_{\text{"big", simple}} \quad (1)$$

with $[H_S, H_B] = 0$.

- ▶ weak coupling $H_I \approx 0$ thermodynamics of open systems are somewhat understood¹
- ▶ strong coupling: $\langle H_I \rangle \sim \langle H_S \rangle \implies$ we can't neglect the interaction \implies thermodynamics?
- ▶ but what is clear: *need to get access to exact dynamics of H_I, H_B*

¹Rivas, "Strong Coupling Thermodynamics of Open Quantum Systems"; Talkner and Hnggi, "Colloquium: Statistical mechanics and thermodynamics at strong coupling: Quantum and classical".

Is that possible?

Is that possible? Yes.

Is that possible? Yes.
Using HOPS :)

Is that possible? Yes.
Using HOPS :)

Sneak Peek

We will be able to calculate $\frac{d\langle H_B \rangle}{dt}$ (and $\langle H_I \rangle$).

▶ more general: $O_S \otimes (B^a)^\dagger B^b$ with $B = \sum_\lambda g_\lambda a_\lambda$

Is that possible? Yes. Using HOPS :)

Sneak Peek

We will be able to calculate $\frac{d\langle H_B \rangle}{dt}$ (and $\langle H_I \rangle$).

▶ more general: $O_S \otimes (B^a)^\dagger B^b$ with $B = \sum_\lambda g_\lambda a_\lambda$

▶ won't call this *heat-flow* because it isn't *the* thermodynamic heat flow

Is that possible? Yes. Using HOPS :)

Sneak Peek

We will be able to calculate $\frac{d\langle H_B \rangle}{dt}$ (and $\langle H_I \rangle$).

▶ more general: $O_S \otimes (B^a)^\dagger B^b$ with $B = \sum_\lambda g_\lambda a_\lambda$

▶ won't call this *heat-flow* because it isn't *the* thermodynamic heat flow

▶ nevertheless: may be interesting *qualitative* measure for energy flow

Introduction

Motivation

Technical Basics

Bath and Interaction Energy

A Little (more) Theory

Analytic Verification

Applications

One Bath

Energy Shovel

Otto Cycle

Anti-Zeno Engine

Outlook

Standard Model of Open Systems

In the following we will work with models of the form²

$$H = H_S(t) + \sum_{n=1}^N \left[H_B^{(n)} + \left(L_n^\dagger(t) B_n + \text{h.c.} \right) \right], \quad (2)$$

where

- ▶ H_S is the System Hamiltonian
- ▶ $H_B^{(n)} = \sum_{\lambda} \omega_{\lambda}^{(n)} a_{\lambda}^{(n),\dagger} a_{\lambda}^{(n)}$
- ▶ $B_n = \sum_{\lambda} g_{\lambda}^{(n)} a_{\lambda}^{(n)}$.

²Sometimes this is called the “Standard Model of Open Systems”.

What remains of the Bath?

Bath Correlation Function

$$\alpha(t-s) = \langle B(t)B(s) \rangle \left(\stackrel{T=0}{=} \sum_{\lambda} |g_{\lambda}|^2 e^{-i\omega_{\lambda}(t-s)} \right) = \frac{1}{\pi} \int J(\omega) e^{-i\omega t} d\omega$$

What remains of the Bath?

Bath Correlation Function

$$\alpha(t-s) = \langle B(t)B(s) \rangle \left(\stackrel{T=0}{=} \sum_{\lambda} |g_{\lambda}|^2 e^{-i\omega_{\lambda}(t-s)} \right) = \frac{1}{\pi} \int J(\omega) e^{-i\omega t} d\omega$$

Spectral Density

$$J(\omega) = \pi \sum_{\lambda} |g_{\lambda}|^2 \delta(\omega - \omega_{\lambda})$$

- ▶ in thermodynamic limit \rightarrow smooth function
- ▶ here usually: Ohmic SD $J(\omega) = \eta\omega e^{-\omega/\omega_c}$ (think phonons)

NMQSD (Zero Temperature)

Open system dynamics formulated as a *stochastic* differential equation:

$$\partial_t \psi_t(\boldsymbol{\eta}_t^*) = -iH(t)\psi_t(\boldsymbol{\eta}_t^*) + \mathbf{L} \cdot \boldsymbol{\eta}_t^* \psi_t(\boldsymbol{\eta}_t^*) - \sum_{n=1}^N L_n^\dagger(t) \int_0^t ds \alpha_n(t-s) \frac{\delta \psi_t(\boldsymbol{\eta}_t^*)}{\delta \eta_n^*(s)}, \quad (3)$$

with

$$\mathcal{M}(\eta_n(t)) = 0, \quad \mathcal{M}(\eta_n(t)\eta_m(s)) = 0, \quad \mathcal{M}(\eta_n(t)\eta_m(s)^*) = \delta_{nm}\alpha_n(t-s), \quad (4)$$

by projecting on coherent bath states.³

³For details see: Disi, Gisin, and W. T. Strunz, “Non-Markovian quantum state diffusion”

HOPS

Using $\alpha_n(\tau) = \sum_{\mu}^{M_n} G_{\mu}^{(n)} e^{-W_{\mu}^{(n)}\tau}$ we define

$$D_{\mu}^{(n)}(t) \equiv \int_0^t ds G_{\mu}^{(n)} e^{-W_{\mu}^{(n)}(t-s)} \frac{\delta}{\delta \eta_n^*(s)} \quad (5)$$

and $D^{\mathbf{k}} \equiv \prod_{n=1}^N \prod_{\mu=1}^{M_n} \sqrt{\frac{\mathbf{k}_{n,\mu}!}{(G_{\mu}^{(n)})^{\mathbf{k}_{n,\mu}}}} \frac{1}{i^{\mathbf{k}_{n,\mu}}} (D_{\mu}^{(n)})^{\mathbf{k}_{n,\mu}}$, $\psi_t^{\mathbf{k}} \equiv D^{\mathbf{k}}\psi_t$ we find

$$\begin{aligned} \dot{\psi}_t^{\mathbf{k}} = & \left[-iH_S(t) + \mathbf{L}(t) \cdot \boldsymbol{\eta}_t^* - \sum_{n=1}^N \sum_{\mu=1}^{M_n} \mathbf{k}_{n,\mu} W_{\mu}^{(n)} \right] \psi_t^{\mathbf{k}} \\ & + i \sum_{n=1}^N \sum_{\mu=1}^{M_n} \sqrt{G_{\mu}^{(n)}} \left[\sqrt{\mathbf{k}_{n,\mu}} L_n(t) \psi_t^{\mathbf{k} - \mathbf{e}_{n,\mu}} + \sqrt{(\mathbf{k}_{n,\mu} + 1)} L_n^{\dagger}(t) \psi_t^{\mathbf{k} + \mathbf{e}_{n,\mu}} \right]. \quad (6) \end{aligned}$$

Introduction

Motivation

Technical Basics

Bath and Interaction Energy

A Little (more) Theory

Analytic Verification

Applications

One Bath

Energy Shovel

Otto Cycle

Anti-Zeno Engine

Outlook

Introduction

Motivation

Technical Basics

Bath and Interaction Energy

A Little (more) Theory

Analytic Verification

Applications

One Bath

Energy Shovel

Otto Cycle

Anti-Zeno Engine

Outlook

Zero Temperature, One Bath, Linear NMQSD

We want to calculate

$$J = -\frac{d\langle H_B \rangle}{dt} = \langle L^\dagger \partial_t B(t) + L \partial_t B^\dagger(t) \rangle_I. \quad (7)$$

Zero Temperature, One Bath, Linear NMQSD

We want to calculate

$$J = -\frac{d\langle H_B \rangle}{dt} = \langle L^\dagger \partial_t B(t) + L \partial_t B^\dagger(t) \rangle_I. \quad (7)$$

...some manipulations ...

Zero Temperature, One Bath, Linear NMQSD

We want to calculate

$$J = -\frac{d\langle H_B \rangle}{dt} = \langle L^\dagger \partial_t B(t) + L \partial_t B^\dagger(t) \rangle_I. \quad (7)$$

...some manipulations ...

Result (NMQSD)

$$J(t) = -i\mathcal{M}_{\eta^*} \langle \psi(\eta, t) | L^\dagger \dot{D}_t | \psi(\eta^*, t) \rangle + \text{c.c.} \quad (8)$$

with $\dot{D}_t = \int_0^t ds \dot{\alpha}(t-s) \frac{\delta}{\delta \eta_s^*}$.

Zero Temperature, One Bath, Linear NMQSD

We want to calculate

$$J = -\frac{d\langle H_B \rangle}{dt} = \langle L^\dagger \partial_t B(t) + L \partial_t B^\dagger(t) \rangle_I. \quad (7)$$

...some manipulations ...

Result (NMQSD)

$$J(t) = -i\mathcal{M}_{\eta^*} \langle \psi(\eta, t) | L^\dagger \dot{D}_t | \psi(\eta^*, t) \rangle + \text{c.c.} \quad (8)$$

with $\dot{D}_t = \int_0^t ds \dot{\alpha}(t-s) \frac{\delta}{\delta \eta_s^*}$.

Result (HOPS)

$$J(t) = -\sum_{\mu} \sqrt{G_{\mu}} W_{\mu} \mathcal{M}_{\eta^*} \langle \psi^{(0)}(\eta, t) | L^\dagger | \psi^{\mathbf{e}_{\mu}}(\eta^*, t) \rangle + \text{c.c.} \quad (9)$$

Generalizations

Finite Temperature

$$J(t) = J_0(t) + [\langle L^\dagger \partial_t \xi(t) \rangle + \text{c.c.}] \quad (10)$$

with $\mathcal{M}(\xi(t)) = 0 = \mathcal{M}(\xi(t)\xi(s))$, $\mathcal{M}(\xi(t)\xi^*(s)) = \frac{1}{\pi} \int_0^\infty d\omega \bar{n}(\beta\omega) J(\omega) e^{-i\omega(t-s)}$ and $J(\omega) = \pi \sum_\lambda |g_\lambda|^2 \delta(\omega - \omega_\lambda)$.⁴

- ▶ nonlinear NMQSD/HOPS
- ▶ multiple baths straight forward
- ▶ interaction energy: “removing the dot”...
- ▶ general “collective” bath observables $O_S \otimes (B^a)^\dagger B^b$ with $B = \sum_\lambda g_\lambda a_\lambda$

⁴ $\partial_t \xi(t)$ exists if correlation function is smooth

Is this any good?

Introduction

Motivation

Technical Basics

Bath and Interaction Energy

A Little (more) Theory

Analytic Verification

Applications

One Bath

Energy Shovel

Otto Cycle

Anti-Zeno Engine

Outlook

Model

$$H = \frac{\Omega}{4}(p^2 + q^2) + \frac{1}{2}q \sum_{\lambda} (g_{\lambda}^* b_{\lambda} + g_{\lambda} b_{\lambda}^{\dagger}) + \sum_{\lambda} \omega_{\lambda} b_{\lambda}^{\dagger} b_{\lambda}, \quad (11)$$

Model

$$H = \frac{\Omega}{4}(p^2 + q^2) + \frac{1}{2}q \sum_{\lambda} (g_{\lambda}^* b_{\lambda} + g_{\lambda} b_{\lambda}^{\dagger}) + \sum_{\lambda} \omega_{\lambda} b_{\lambda}^{\dagger} b_{\lambda}, \quad (11)$$

...leading to ...

$$\dot{q} = \Omega p \quad (12)$$

$$\dot{p} = -\Omega q - \int_0^t \mathfrak{I}[\alpha_0(t-s)]q(s) ds + W(t) \quad (13)$$

$$\dot{b}_{\lambda} = -ig_{\lambda} \frac{q}{2} - i\omega_{\lambda} b_{\lambda} \quad (14)$$

with the operator noise $W(t) = -\sum_{\lambda} (g_{\lambda}^* b_{\lambda}(0)e^{-i\omega_{\lambda}t} + g_{\lambda} b_{\lambda}^{\dagger}(0)e^{i\omega_{\lambda}t})$,
 $\langle W(t)W(s) \rangle = \alpha(t-s)$ and $\alpha_0 \equiv \alpha|_{T=0}$.

Solution through a matrix $G(t)$ with $G(0) = \mathbb{1}$ and

$$\dot{G}(t) = AG(t) - \int_0^t K(t-s)G(s) \, ds, \quad A = \begin{pmatrix} 0 & \Omega \\ -\Omega & 0 \end{pmatrix}, \quad K(t) = \begin{pmatrix} 0 & 0 \\ \mathfrak{I}[\alpha_0(t)] & 0 \end{pmatrix}. \quad (15)$$

Solution through a matrix $G(t)$ with $G(0) = \mathbb{1}$ and

$$\dot{G}(t) = AG(t) - \int_0^t K(t-s)G(s) \, ds, \quad A = \begin{pmatrix} 0 & \Omega \\ -\Omega & 0 \end{pmatrix}, \quad K(t) = \begin{pmatrix} 0 & 0 \\ \mathfrak{I}[\alpha_0(t)] & 0 \end{pmatrix}. \quad (15)$$

Then

$$\begin{pmatrix} q(t) \\ p(t) \end{pmatrix} = G(t) \begin{pmatrix} q(0) \\ p(0) \end{pmatrix} + \int_0^t G(t-s) \begin{pmatrix} 0 \\ W(s) \end{pmatrix} \, ds. \quad (16)$$

► “exact” solution via laplace transform and BCF expansion + residue theorem

Result

Solution

$$G(t) = \sum_{l=1}^{N+1} \left[R_l \begin{pmatrix} \tilde{z}_l & \Omega \\ \frac{\tilde{z}_l^2}{\Omega} & \tilde{z}_l \end{pmatrix} e^{\tilde{z}_l \cdot t} + \text{c.c.} \right] \quad (17)$$

with $R_l = f_0(\tilde{z}_l)/p'(\tilde{z}_l)$, f_0, p polynomials, \tilde{z}_l roots of p .

Result

Solution

$$G(t) = \sum_{l=1}^{N+1} \left[R_l \begin{pmatrix} \tilde{z}_l & \Omega \\ \frac{\tilde{z}_l^2}{\Omega} & \tilde{z}_l \end{pmatrix} e^{\tilde{z}_l \cdot t} + \text{c.c.} \right] \quad (17)$$

with $R_l = f_0(\tilde{z}_l)/p'(\tilde{z}_l)$, f_0, p polynomials, \tilde{z}_l roots of p .

- ▶ note: G doesn't depend on temperature
- ▶ solution very sensitive to precision of the fits and roots

Bath Energy Derivative



$$\begin{aligned}\langle \dot{H}_B \rangle &= \sum_{\lambda} \omega_{\lambda} (\langle b_{\lambda}^{\dagger} \dot{b}_{\lambda} \rangle + \text{c.c.}) \\ &= -\frac{1}{2} \Im \left[\int_0^t ds \langle q(t) q(s) \rangle \dot{\alpha}_0(t-s) \right] \\ &\quad + \frac{1}{2} G_{12}(t) [\alpha(t) - \alpha_0(t)] - \frac{\Omega}{2} \int_0^t ds G_{11}(s) [\alpha(s) - \alpha_0(s)]\end{aligned}\tag{18}$$

► becomes huge sum of exponentials (thanks Mathematica)

One Bath, Finite Temperature

Parameters



$\Omega = 1$, Ohmic BCF $\frac{\eta}{\pi}(\omega_c/(1+i\omega_c\tau))^2$ with $(\alpha(0) = 0.64, \omega_c = 2)$, $N = 10^5$ samples, 15 Hilbert space dimensions, $|\psi(0)\rangle_S = |1\rangle_S$, $T = 1$



Two Baths, Finite Temperature (Gradient)

Parameters

$\Omega = \Lambda = 1$, symmetric Ohmic BCFs with $(\alpha(0) = 0.25, \omega_c = 2)$, $N = 10^4$ samples, 9 Hilbert space dimensions, $|\psi(0)\rangle_S = |0, 0\rangle_S$, $T = 0.6$, $\gamma = 0.5$



Introduction

Motivation

Technical Basics

Bath and Interaction Energy

A Little (more) Theory

Analytic Verification

Applications

One Bath

Energy Shovel

Otto Cycle

Anti-Zeno Engine

Outlook

Introduction

Motivation

Technical Basics

Bath and Interaction Energy

A Little (more) Theory

Analytic Verification

Applications

One Bath

Energy Shovel

Otto Cycle

Anti-Zeno Engine

Outlook

One Bath, Zero Temperature

Model: Spin-Boson

$$H = \frac{1}{2}\sigma_z + \frac{1}{2} \sum_{\lambda} (g_{\lambda}\sigma_x^{\dagger}a_{\lambda} + g_{\lambda}^*\sigma_x a_{\lambda}^{\dagger}) + \sum_{\lambda} \omega_{\lambda} a_{\lambda}^{\dagger} a_{\lambda}, |\psi_0\rangle_S = |\uparrow\rangle \quad (19)$$

► how do we check convergence:

One Bath, Zero Temperature

Model: Spin-Boson

$$H = \frac{1}{2}\sigma_z + \frac{1}{2} \sum_{\lambda} (g_{\lambda}\sigma_x^{\dagger}a_{\lambda} + g_{\lambda}^*\sigma_x a_{\lambda}^{\dagger}) + \sum_{\lambda} \omega_{\lambda} a_{\lambda}^{\dagger} a_{\lambda}, |\psi_0\rangle_S = |\uparrow\rangle \quad (19)$$

► how do we check convergence:

One Bath, Zero Temperature

Model: Spin-Boson

$$H = \frac{1}{2}\sigma_z + \frac{1}{2} \sum_{\lambda} (g_{\lambda}\sigma_x^{\dagger}a_{\lambda} + g_{\lambda}^*\sigma_x a_{\lambda}^{\dagger}) + \sum_{\lambda} \omega_{\lambda} a_{\lambda}^{\dagger} a_{\lambda}, |\psi_0\rangle_S = |\uparrow\rangle \quad (19)$$

- ▶ how do we check convergence:
 - ▶ old: difference of results to some “good” configuration


One Bath, Zero Temperature

Model: Spin-Boson

$$H = \frac{1}{2}\sigma_z + \frac{1}{2} \sum_{\lambda} (g_{\lambda}\sigma_x^{\dagger}a_{\lambda} + g_{\lambda}^*\sigma_x a_{\lambda}^{\dagger}) + \sum_{\lambda} \omega_{\lambda} a_{\lambda}^{\dagger} a_{\lambda}, |\psi_0\rangle_S = |\uparrow\rangle \quad (19)$$

- ▶ how do we check convergence:
 - ▶ old: difference of results to some “good” configuration
 - ▶ new: consistency with energy conservation





Example: Dependence of the Interaction Energy on Stochastic Process Sampling






- ▶ $\alpha(0) = 1.6$ and $\omega_c = 4 \implies$ stress HOPS through fast decaying BCF
- ▶ “perfect” results only with very high accuracy⁵ ζ
- ▶ good qualitative results for less extreme configurations (common theme)

⁵smaller ζ is better

Various Cutoff Frequencies





Non-Markovian Dynamics



► interaction strengths chosen for approx. same interaction energy

Non-Markovian Dynamics



- ▶ interaction strengths chosen for approx. same interaction energy
- ▶ timing important for energy transfer “performance”

Beware :)

The following is WIP and has not been written up properly yet.

Introduction

Motivation

Technical Basics

Bath and Interaction Energy

A Little (more) Theory

Analytic Verification

Applications

One Bath

Energy Shovel


Otto Cycle

Anti-Zeno Engine

Outlook


Extracting Energy from One Bath

- ▶ same model as above eq. (19), but with $L(\tau) = \sin^2(\frac{\Delta}{2}\tau)\sigma_x$
- ▶ how much energy can be *unitarily* extracted? $\implies \Delta E_{\max} = \frac{1}{\beta} S(\rho_S \parallel \rho_S^\beta)$



Extracting Energy from One Bath

- ▶ same model as above eq. (19), but with $L(\tau) = \sin^2(\frac{\Delta}{2}\tau)\sigma_x$
- ▶ how much energy can be *unitarily* extracted? $\implies \Delta E_{\max} = \frac{1}{\beta} S(\rho_S \parallel \rho_S^\beta)$



Introduction

Motivation

Technical Basics

Bath and Interaction Energy

A Little (more) Theory

Analytic Verification

Applications

One Bath

Energy Shovel

Otto Cycle

Anti-Zeno Engine


Outlook

Otto Cycle (proof of concept)


Model


Spin-Boson model with compression of H_S and modulation of L .


► classical toy model of the quantum heat engine community⁶



⁶Geva and Kosloff, "A quantum-mechanical heat engine operating in finite time. A model consisting of spin-1/2 systems as the working fluid".







- ▶ $\bar{P} = 1.98 \cdot 10^{-3} \pm 2.3 \cdot 10^{-5}$, $\eta \approx 20\%$, $T_c = 1$, $T_h = 20$
- ▶ no tuning of parameters, except for resonant coupling
- ▶ long bath memory $\omega_c = 1$, but weak-ish coupling

Questions (for the future)

- ▶ better performance through “overlapping” phases?
- ▶ strong coupling any good?
- ▶ non-Markovianity + strong coupling any good?
- ▶ what is the optimal efficiency and power? (probably no advantage here)

Introduction

Motivation

Technical Basics

Bath and Interaction Energy

A Little (more) Theory

Analytic Verification

Applications

One Bath

Energy Shovel

Otto Cycle

Anti-Zeno Engine

Outlook

Anti-Zeno Engine

Question

Is there a use for non-Markovianity in quantum heat engines?


- ▶ Mukherjee, Kofman, and Kurizki, “Anti-Zeno quantum advantage in fast-driven heat machines” claims that one can exploit the time-energy uncertainty for quantum advantage⁷

Model


Qubit coupled to two baths of different temperatures (T_c, T_h)

$$H_S = \frac{1}{2}[\omega_0 + \gamma\Delta \sin(\Delta t)]\sigma_z, \quad L_{c,h} = \frac{1}{2}\sigma_x \quad (20)$$


⁷I'd be careful to call this quantum advantage.



- ▶ couple for n modulation periods slightly of resonance
- ▶ for smaller n the $\frac{\sin((\omega - (\omega_0 \pm \Delta))\tau)}{((\omega - (\omega_0 \pm \Delta))\tau)}$ has a greater overlap \implies controls power output



a) $P = -0.058 \pm 0.014$



b) $P = -0.068 \pm 0.010$

Parameters

$$\Delta = 11, \gamma = 0.5, \alpha(0) = 1.0, \omega_0 = 20, T_c = 8, T_h = 40$$

- ▶ this is not properly converged yet \rightarrow newer results: no advantage at these temperatures / coupling strengths
- ▶ new simulations with temperatures from paper ($\beta_{h(c)} = 0.0005(0.005)$) are promising
 - ▶ interesting \rightarrow no good steady state power in this case (insufficient samples?)

Introduction

Motivation

Technical Basics

Bath and Interaction Energy

A Little (more) Theory

Analytic Verification

Applications

One Bath

Energy Shovel

Otto Cycle

Anti-Zeno Engine

Outlook






On the “To Do” List

- ▶ verify/falsify weak coupling results in the literature (engines)
- ▶ three-level systems: there is an experimental paper ;)
- ▶ parameter scan of two qubit model
- ▶ filter modes
- ▶ ...







Lessons Learned

- ▶ careful convergence checks pay off
- ▶ surveying literature is important
- ▶ properly documenting observations is a great help and should be done as early as possible
- ▶ applications should be carefully chosen to answer interesting questions
- ▶ numerics are helpful, but physical insights are important
- ▶ comparison with some experiments would have been nice







References I

-  Bera, Mohit Lal, Sergi Juli-Farr, et al. “Quantum Heat Engines with Carnot Efficiency at Maximum Power”. In: *arXiv* (June 2021). eprint: 2106.01193. URL: <https://arxiv.org/abs/2106.01193v1>.
-  Bera, Mohit Lal, Maciej Lewenstein, and Manabendra Nath Bera. “Attaining Carnot efficiency with quantum and nanoscale heat engines - npj Quantum Information”. In: *npj Quantum Inf.* 7.31 (Feb. 2021), pp. 1–7. ISSN: 2056-6387. DOI: 10.1038/s41534-021-00366-6.
-  Disi, L., N. Gisin, and W. T. Strunz. “Non-Markovian quantum state diffusion”. In: *Phys. Rev. A* 58.3 (Sept. 1998), pp. 1699–1712. ISSN: 2469-9934. DOI: 10.1103/PhysRevA.58.1699.
-  Esposito, Massimiliano, Maicol A. Ochoa, and Michael Galperin. “Nature of heat in strongly coupled open quantum systems”. In: *Phys. Rev. B* 92.23 (Dec. 2015), p. 235440. ISSN: 2469-9969. DOI: 10.1103/PhysRevB.92.235440.
-  Gao, Xing et al. “Non-Markovian Stochastic Schrödinger Equation: Matrix Product State Approach to the Hierarchy of Pure States”. In: *arXiv* (Sept. 2021). eprint: 2109.06393. URL: <https://arxiv.org/abs/2109.06393v3>.


References II

-  Geva, Eitan and Ronnie Kosloff. “A quantum-mechanical heat engine operating in finite time. A model consisting of spin-1/2 systems as the working fluid”. In: *J. Chem. Phys.* 96.4 (Feb. 1992), pp. 3054–3067. ISSN: 0021-9606. DOI: 10.1063/1.461951.
-  Kato, Akihito and Yoshitaka Tanimura. “Quantum heat current under non-perturbative and non-Markovian conditions: Applications to heat machines”. In: *J. Chem. Phys.* 145.22 (Dec. 2016), p. 224105. ISSN: 0021-9606. DOI: 10.1063/1.4971370.
-  — . “Quantum heat transport of a two-qubit system: Interplay between system-bath coherence and qubit-qubit coherence”. In: *J. Chem. Phys.* 143.6 (Aug. 2015), p. 064107. ISSN: 0021-9606. DOI: 10.1063/1.4928192.
-  Klauder, JR and ECG Sudarshan. “Fundamentals of Quantum Optics Benjamin”. In: *Inc., New York* (1968).
-  Motz, T. et al. “Rectification of heat currents across nonlinear quantum chains: a versatile approach beyond weak thermal contact”. In: *New J. Phys.* 20.11 (Nov. 2018), p. 113020. ISSN: 1367-2630. DOI: 10.1088/1367-2630/aaea90.
-  Mukherjee, Victor, Abraham G. Kofman, and Gershon Kurizki. “Anti-Zeno quantum advantage in fast-driven heat machines”. In: *Commun. Phys.* 3.8 (Jan. 2020), pp. 1–12. ISSN: 2399-3650. DOI: 10.1038/s42005-019-0272-z.

References III

-  Rivas, ngel. “Strong Coupling Thermodynamics of Open Quantum Systems”. In: *arXiv* (Oct. 2019). DOI: [10.1103/PhysRevLett.124.160601](https://doi.org/10.1103/PhysRevLett.124.160601). eprint: [1910.01246](https://arxiv.org/abs/1910.01246).
-  Senior, Jorden et al. “Heat rectification via a superconducting artificial atom - Communications Physics”. In: *Commun. Phys.* 3.40 (Feb. 2020), pp. 1–5. ISSN: 2399-3650. DOI: [10.1038/s42005-020-0307-5](https://doi.org/10.1038/s42005-020-0307-5).
-  Strasberg, Philipp and Andreas Winter. “First and Second Law of Quantum Thermodynamics: A Consistent Derivation Based on a Microscopic Definition of Entropy”. In: *PRX Quantum* 2.3 (Aug. 2021), p. 030202. ISSN: 2691-3399. DOI: [10.1103/PRXQuantum.2.030202](https://doi.org/10.1103/PRXQuantum.2.030202).
-  Strunz, Walter T. “Stochastic Schrödinger equation approach to the dynamics of non-Markovian open quantum systems”. *Fachbereich Physik der Universität Essen*, 2001.
-  Talkner, Peter and Peter Hnggi. “Colloquium: Statistical mechanics and thermodynamics at strong coupling: Quantum and classical”. In: *Rev. Mod. Phys.* 92.4 (Oct. 2020), p. 041002. ISSN: 1539-0756. DOI: [10.1103/RevModPhys.92.041002](https://doi.org/10.1103/RevModPhys.92.041002).
-  — . “Open system trajectories specify fluctuating work but not heat”. In: *Phys. Rev. E* 94.2 (Aug. 2016), p. 022143. ISSN: 2470-0053. DOI: [10.1103/PhysRevE.94.022143](https://doi.org/10.1103/PhysRevE.94.022143).

References IV

-  Wiedmann, M., J. T. Stockburger, and J. Ankerhold. “Non-Markovian dynamics of a quantum heat engine: out-of-equilibrium operation and thermal coupling control”. In: *New J. Phys.* 22.3 (Mar. 2020), p. 033007. ISSN: 1367-2630. DOI: [10.1088/1367-2630/ab725a](https://doi.org/10.1088/1367-2630/ab725a).

Situation (Longer)

Consider an open quantum system

$$H = \underbrace{H_S}_{\text{"small"}} + \underbrace{H_I}_{?} + \underbrace{H_B}_{\text{"big", simple}} \quad (21)$$

with $[H_S, H_B] = 0$.

⁸even in strong coupling equilibrium...

⁹Rivas, "Strong Coupling Thermodynamics of Open Quantum Systems"; Talkner and Hnggi, "Colloquium: Statistical mechanics and thermodynamics at strong coupling: Quantum and classical".

Situation (Longer)

Consider an open quantum system

$$H = \underbrace{H_S}_{\text{"small"}} + \underbrace{H_I}_{?} + \underbrace{H_B}_{\text{"big", simple}} \quad (21)$$

with $[H_S, H_B] = 0$.

- ▶ weak coupling $H_I \approx 0$ thermodynamics⁸ of open systems are somewhat understood⁹

⁸even in strong coupling equilibrium...

⁹Rivas, "Strong Coupling Thermodynamics of Open Quantum Systems"; Talkner and Hnggi, "Colloquium: Statistical mechanics and thermodynamics at strong coupling: Quantum and classical".

Situation (Longer)

Consider an open quantum system

$$H = \underbrace{H_S}_{\text{"small"}} + \underbrace{H_I}_{?} + \underbrace{H_B}_{\text{"big", simple}} \quad (21)$$

with $[H_S, H_B] = 0$.

- ▶ weak coupling $H_I \approx 0$ thermodynamics⁸ of open systems are somewhat understood⁹
- ▶ strong coupling: $\langle H_I \rangle \sim \langle H_S \rangle \implies$ we can't neglect the interaction \implies thermodynamics?

⁸even in strong coupling equilibrium...

⁹Rivas, "Strong Coupling Thermodynamics of Open Quantum Systems"; Talkner and Hnggi, "Colloquium: Statistical mechanics and thermodynamics at strong coupling: Quantum and classical".

Situation (Longer)

Consider an open quantum system

$$H = \underbrace{H_S}_{\text{"small"}} + \underbrace{H_I}_{?} + \underbrace{H_B}_{\text{"big", simple}} \quad (21)$$

with $[H_S, H_B] = 0$.

- ▶ weak coupling $H_I \approx 0$ thermodynamics⁸ of open systems are somewhat understood⁹
- ▶ strong coupling: $\langle H_I \rangle \sim \langle H_S \rangle \implies$ we can't neglect the interaction \implies thermodynamics?
- ▶ we do quantum mechanics \implies can't separate bath and system

⁸even in strong coupling equilibrium...

⁹Rivas, "Strong Coupling Thermodynamics of Open Quantum Systems"; Talkner and Hnggi, "Colloquium: Statistical mechanics and thermodynamics at strong coupling: Quantum and classical".

Situation (Longer)

Consider an open quantum system

$$H = \underbrace{H_S}_{\text{"small"}} + \underbrace{H_I}_{?} + \underbrace{H_B}_{\text{"big", simple}} \quad (21)$$

with $[H_S, H_B] = 0$.

- ▶ weak coupling $H_I \approx 0$ thermodynamics⁸ of open systems are somewhat understood⁹
- ▶ strong coupling: $\langle H_I \rangle \sim \langle H_S \rangle \implies$ we can't neglect the interaction \implies thermodynamics?
- ▶ we do quantum mechanics \implies can't separate bath and system, especially not dynamics!

⁸even in strong coupling equilibrium...

⁹Rivas, "Strong Coupling Thermodynamics of Open Quantum Systems"; Talkner and Hnggi, "Colloquium: Statistical mechanics and thermodynamics at strong coupling: Quantum and classical".

Situation (Longer)

Consider an open quantum system

$$H = \underbrace{H_S}_{\text{"small"}} + \underbrace{H_I}_{?} + \underbrace{H_B}_{\text{"big", simple}} \quad (21)$$

with $[H_S, H_B] = 0$.

- ▶ weak coupling $H_I \approx 0$ thermodynamics⁸ of open systems are somewhat understood⁹
- ▶ strong coupling: $\langle H_I \rangle \sim \langle H_S \rangle \implies$ we can't neglect the interaction \implies thermodynamics?
- ▶ we do quantum mechanics \implies can't separate bath and system, especially not dynamics!
- ▶ no consensus about strong coupling thermodynamics:

⁸even in strong coupling equilibrium...

⁹Rivas, "Strong Coupling Thermodynamics of Open Quantum Systems"; Talkner and Hnggi, "Colloquium: Statistical mechanics and thermodynamics at strong coupling: Quantum and classical".

Situation (Longer)

Consider an open quantum system

$$H = \underbrace{H_S}_{\text{"small"}} + \underbrace{H_I}_{?} + \underbrace{H_B}_{\text{"big", simple}} \quad (21)$$

with $[H_S, H_B] = 0$.

- ▶ weak coupling $H_I \approx 0$ thermodynamics⁸ of open systems are somewhat understood⁹
- ▶ strong coupling: $\langle H_I \rangle \sim \langle H_S \rangle \implies$ we can't neglect the interaction \implies thermodynamics?
- ▶ we do quantum mechanics \implies can't separate bath and system, especially not dynamics!
- ▶ no consensus about strong coupling thermodynamics:
- ▶ but what is clear: *need to get access to exact dynamics of H_I, H_B*

⁸even in strong coupling equilibrium...


⁹Rivas, "Strong Coupling Thermodynamics of Open Quantum Systems"; Talkner and Hnggi, "Colloquium: Statistical mechanics and thermodynamics at strong coupling: Quantum and classical".

More Papers on Thermo


M. L. Bera, Juli-Farr, et al., “Quantum Heat Engines with Carnot Efficiency at Maximum Power”; M. L. Bera, Lewenstein, and M. N. Bera, “Attaining Carnot efficiency with quantum and nanoscale heat engines - npj Quantum Information”; Esposito, Ochoa, and Galperin, “Nature of heat in strongly coupled open quantum systems”; Kato and Tanimura, “Quantum heat current under non-perturbative and non-Markovian conditions: Applications to heat machines”, “Quantum heat transport of a two-qubit system: Interplay between system-bath coherence and qubit-qubit coherence”; Motz et al., “Rectification of heat currents across nonlinear quantum chains: a versatile approach beyond weak thermal contact”; Rivas, “Strong Coupling Thermodynamics of Open Quantum Systems”; Senior et al., “Heat rectification via a superconducting artificial atom - Communications Physics”; Strasberg and Winter, “First and Second Law of Quantum Thermodynamics: A Consistent Derivation Based on a Microscopic Definition of Entropy”; Talkner and Hnggi, “Colloquium: Statistical mechanics and thermodynamics at strong coupling: Quantum and classical”, “Open system trajectories specify fluctuating work but not heat”; Wiedmann, Stockburger, and Ankerhold, “Non-Markovian dynamics of a quantum heat engine: out-of-equilibrium operation and thermal coupling control”

Ohmic SD BCF

$$J = \eta e^{-\omega/\omega_c} \omega^s$$



$$J = \eta \Gamma(s+1) / (1 + i\omega_c \tau)^{s+1}$$



NMQSD (Zero Temperature)

Expanding in a Bargmann (unnormalized) coherent state basis Klauder and Sudarshan, “Fundamentals of Quantum Optics Benjamin” $\{|\mathbf{z}^{(1)}, \mathbf{z}^{(2)}, \dots\rangle = |\underline{\mathbf{z}}\rangle\}$

$$|\psi(t)\rangle = \int \prod_{n=1}^N \left(\frac{d\mathbf{z}^{(n)}}{\pi^{N_n}} e^{-|\mathbf{z}|^2} \right) |\psi(t, \underline{\mathbf{z}}^*)\rangle |\underline{\mathbf{z}}\rangle, \quad (22)$$

we obtain

$$\partial_t \psi_t(\boldsymbol{\eta}_t^*) = -iH\psi_t(\boldsymbol{\eta}_t^*) + \mathbf{L} \cdot \boldsymbol{\eta}_t^* \psi_t(\boldsymbol{\eta}_t^*) - \sum_{n=1}^N L_n^\dagger \int_0^t ds \alpha_n(t-s) \frac{\delta \psi_t(\boldsymbol{\eta}_t^*)}{\delta \eta_n^*(s)}, \quad (23)$$

with

$$\mathcal{M}(\eta_n^*(t)) = 0, \quad \mathcal{M}(\eta_n(t)\eta_m(s)) = 0, \quad \mathcal{M}(\eta_n(t)\eta_m(s)^*) = \delta_{nm} \alpha_n(t-s), \quad (24)$$

where $\alpha_n(t-s) = \sum_\lambda |g_\lambda^{(n)}|^2 e^{-i\omega_\lambda^{(n)}(t-s)} = \langle B(t)B(s) \rangle_{I, \rho(0)}$ Walter T. Strunz, “Stochastic Schrödinger equation approach to the dynamics of non-Markovian open quantum systems” (fourier transf. of spectral density $J(\omega) = \pi \sum_\lambda |g_\lambda|^2 \delta(\omega - \omega_\lambda)$).

Fock-Space Embedding

As in Gao et al., “Non-Markovian Stochastic Schrödinger Equation: Matrix Product State Approach to the Hierarchy of Pure States” we can define

$$|\Psi\rangle = \sum_{\underline{\mathbf{k}}} |\psi^{\underline{\mathbf{k}}}\rangle \otimes |\underline{\mathbf{k}}\rangle \quad (25)$$

where $|\underline{\mathbf{k}}\rangle = \bigotimes_{n=1}^N \bigotimes_{\mu=1}^{M_n} |\mathbf{k}_{n,\mu}\rangle$ are bosonic Fock-states.

Now eq. (6) becomes

$$\partial_t |\Psi\rangle = \left[-iH_S + \mathbf{L} \cdot \boldsymbol{\eta}^* - \sum_{n=1}^N \sum_{\mu=1}^{M_n} b_{n,\mu}^\dagger b_{n,\mu} W_\mu^{(n)} + i \sum_{n=1}^N \sum_{\mu=1}^{M_n} \sqrt{G_{n,\mu}} (b_{n,\mu}^\dagger L_n + b_{n,\mu} L_n^\dagger) \right] |\Psi\rangle. \quad (26)$$

Fock-Space Embedding

As in Gao et al., “Non-Markovian Stochastic Schrödinger Equation: Matrix Product State Approach to the Hierarchy of Pure States” we can define

$$|\Psi\rangle = \sum_{\underline{\mathbf{k}}} |\psi^{\underline{\mathbf{k}}}\rangle \otimes |\underline{\mathbf{k}}\rangle \quad (25)$$

where $|\underline{\mathbf{k}}\rangle = \bigotimes_{n=1}^N \bigotimes_{\mu=1}^{M_n} |\underline{\mathbf{k}}_{n,\mu}\rangle$ are bosonic Fock-states.

Now eq. (6) becomes

$$\partial_t |\Psi\rangle = \left[-iH_S + \mathbf{L} \cdot \boldsymbol{\eta}^* - \sum_{n=1}^N \sum_{\mu=1}^{M_n} b_{n,\mu}^\dagger b_{n,\mu} W_\mu^{(n)} + i \sum_{n=1}^N \sum_{\mu=1}^{M_n} \sqrt{G_{n,\mu}} (b_{n,\mu}^\dagger L_n + b_{n,\mu} L_n^\dagger) \right] |\Psi\rangle. \quad (26)$$

\implies possible to derive an upper bound for the norm of $|\psi^{\underline{\mathbf{k}}}\rangle$

Fock-Space Embedding

As in Gao et al., “Non-Markovian Stochastic Schrödinger Equation: Matrix Product State Approach to the Hierarchy of Pure States” we can define

$$|\Psi\rangle = \sum_{\underline{\mathbf{k}}} |\psi^{\underline{\mathbf{k}}}\rangle \otimes |\underline{\mathbf{k}}\rangle \quad (25)$$

where $|\underline{\mathbf{k}}\rangle = \bigotimes_{n=1}^N \bigotimes_{\mu=1}^{M_n} |\underline{\mathbf{k}}_{n,\mu}\rangle$ are bosonic Fock-states.

Now eq. (6) becomes

$$\partial_t |\Psi\rangle = \left[-iH_S + \mathbf{L} \cdot \boldsymbol{\eta}^* - \sum_{n=1}^N \sum_{\mu=1}^{M_n} b_{n,\mu}^\dagger b_{n,\mu} W_\mu^{(n)} + i \sum_{n=1}^N \sum_{\mu=1}^{M_n} \sqrt{G_{n,\mu}} (b_{n,\mu}^\dagger L_n + b_{n,\mu} L_n^\dagger) \right] |\Psi\rangle. \quad (26)$$

\Rightarrow possible to derive an upper bound for the norm of $|\psi^{\underline{\mathbf{k}}}\rangle \Rightarrow$ new truncation scheme